

**Research Article**

DOI : 10.15740/HAS/AJSS/11.1/207-212

# Co-inoculation of phosphate solubilizing bacteria and *Rhizobia* for improving growth and yield of mungbean (*Vigna radiata* L.)

■ H.K. TARAFDER, A. DEY AND S. DASGUPTA

Received : 16.02.2016; Revised : 22.04.2016; Accepted : 18.05.2016

**MEMBERS OF RESEARCH FORUM:**

**Corresponding author :**

**H. K. TARAFDER**, Regional Research Station (U.B.K.V.) (Hill Zone) Kalimpong, DARJEELING (W.B.) INDIA  
Email: [hridaykamalt25@gmail.com](mailto:hridaykamalt25@gmail.com)

**Co-authors :**

**A. DEY**, Regional Research Station (B.C.K.V.), New Alluvial Zone, Gayeshpur, NADIA (W.B.) INDIA  
Email: [arupdey89@gmail.com](mailto:arupdey89@gmail.com)

**S. DASGUPTA**, AICRP on STCR, Directorate of Reserach, Bidhan Chandra Krishi Viswavidyalaya, Kalyani, NADIA (W.B.) INDIA  
Email: [sdasgupta173@gmail.com](mailto:sdasgupta173@gmail.com)

**Summary**

Mung bean is an important pulse crop in West Bengal, growing nearly 11.7 thousands hectares of land. Scientific literature and case studies of inoculating of microbes by various scientists found significant response to crop growth. The present study was made to reduce fertilizer application rate by coinnoculating phosphate solubilizing bacteria and rhizobia for mung bean. The experiment was conducted in the year 2011 and 2012 at the 'Instructional Farm' Bidhan Chandra Krishi Viswavidyalaya, Mohanpur, Nadia, West Bengal. Factorial Randomized Block Design was laid out with three replications and 10 treatment combinations. The treatments was without inoculation ( $A_0$ ) and seed inoculation with *Rhizobium* and phosphate solubilizing bacteria (PSB) strain ( $A_1$ ) as factor one and fertilizer treatments like untreated control (no fertilizer application  $B_0$ ), application of NPK (recommended dose as basal  $B_1$ ), 75 per cent recommended dose of N and  $P_2O_5 + 100\% K_2O$  ( $B_2$ ) and 50 per cent recommended dose of N and  $P_2O_5 + 100\% K_2O$  ( $B_3$ ). It was observed that inoculation with *Rhizobium* and phosphate solubilizing bacteria (PSB) along with 75 per cent RDF *i.e.* treatment combination ( $A_1B_2$ ) was at par with treatment  $A_1B_1$  *i.e.* 100 per cent RDF with respect to all the growth parameters and yield attributing characters of mung bean. So, it can be concluded that both 25 per cent nitrogenous and phosphatic fertilizer of the recommended dose can be substituted by seed co-inoculation with phosphate solubilizing bacteria and rhizobia without affecting the yield compared to 100 per cent RDF.

**Key words :** *Bacillus polymyxa*, *Rhizobia*, Mung bean, Seed inoculation

**How to cite this article :** Tarafder, H.K., Dey, A. and Dasgupta, S. (2016). Co-inoculation of phosphate solubilizing bacteria and *Rhizobia* for improving growth and yield of mungbean (*Vigna radiata* L.). *Asian J. Soil Sci.*, 11 (1) : 207-212 : DOI : 10.15740/HAS/AJSS/11.1/207-212.

## Introduction

Govt. of India now focusing to increase area, production and productivity of pulse crop. Pulse growing area in India nearly 23 million ha with the annual production of 14.18 million tonnes and an average productivity of 617 kg ha<sup>-1</sup> (GOI, 2009). Even being the

largest producer in the World, our country has to import pulses about two million tones every year to feed our nation (Chaturvedi and Ali, 2002). Among the pulse crop mung bean [*Vigna radiata* (L.) Wilczek], commonly known as 'Moong or Mung bean'; is a good source of high quality protein (25%) with high digestibility (Payasi

*et al.*, 2011). In West Bengal, about 4.4 thousand tonnes of mungbean production was recorded from 11.7 thousand hectares of area in 2005-06 (GOWB, 2006). It is well known that Mung bean being leguminous plant form symbiotic association by *Rhizobium* and thereby fixes atmospheric nitrogen (Singh *et al.*, 2011) and reduces nitrogen fertilizer requirement. Phosphorus is vital component of the substances that are building block of genes and chromosomes and the second major primary nutrient limiting plant growth is generally deficient in most of the soils due to its ready fixation (Schachtman *et al.*, 1998). The increasing prices of phosphatic fertilizers have raised an alarming situation for our country. Therefore, it is very tricky for poor farmers to supplement P fertilizers in the soil to circumvent the P deficiency. The combined application of phosphate solubilizing bacteria and nodule forming bacteria in legumes stimulated plant growth (Algawadi and Gaur, 1988 and Perveen *et al.*, 2002) as compared to their single inoculation.

Co-inoculation with N<sub>2</sub>-fixing and P solubilizing microbe was more effective approach for providing balanced plant nutrition (Gupta *et al.*, 1998 and Martins *et al.*, 2004). Present study was designed to evaluate the co-inoculation potential of *Bacillus polymyxa* and *Rhizobium strain* (M-10) for improving growth, nodulation and yield of mung bean at different fertilizer levels.

## Resource and Research Methods

The experiment was conducted in the month between March to May in the year 2011 and 2012 at the 'Instructional Farm' (22°95' north latitude, 88°50' East longitude), Bidhan Chandra Krishi Viswavidyalaya, Mohanpur, Nadia, West Bengal. Soil was sandy-loam in nature, neutral in reaction pH 6.5, with available N (Subbiah and Asija, 1956), P (Olsen *et al.*, 1954) and K (Brown and Wranckle, 1988) was 178.7, 17.72 and 157.52.0 kg ha<sup>-1</sup>, respectively. Factorial Randomized Block Design was laid out with three replications and 10 treatment combinations. The treatments was without inoculation (A<sub>0</sub>) and seed inoculation with *Rhizobium* and phosphate solubilizing bacteria (PSB) strain (A<sub>1</sub>) as factor one and fertilizer treatments like untreated control (no fertilizer application B<sub>0</sub>), application of NPK (recommended dose as basal B<sub>1</sub>), 75 per cent recommended dose of N and P<sub>2</sub>O<sub>5</sub> + 100% K<sub>2</sub>O (B<sub>2</sub>) and 50 per cent recommended dose of N and P<sub>2</sub>O<sub>5</sub> + 100% K<sub>2</sub>O (B<sub>3</sub>).

The *Rhizobium* strains (M-10) and phosphate solubilizing bacteria (*Bacillus polymyxa*) culture was collected from Nodule Research Laboratory, B.C.K.V. For A<sub>1</sub> treatment seeds were mixed thoroughly with the *Rhizobium* and phosphate solubilizing bacteria (*Bacillus polymyxa*) followed by shade drying for 30 minutes before sowing. In case of fertilizer treatment fertilizers were applied as basal. The green gram crop (cv. B-1) was sown at plant to plant spacing of 10 cm and row to row spacing of 30 cm apart. Other agronomic practices done as usual for pulse crop. Growth parameters like plant height, dry matter accumulation, crop growth rate, number and dry weight of nodules were recorded at different growth stages of crop and yield attributing characters and yield were noted at the time of harvest. The experiment was analyzed following the standard method of statistical analysis for factorial RBD.

## Research Findings and Discussion

The results obtained from the present investigation as well as relevant discussion have been summarized under following heads :

### Effect on growth attributes :

Inoculation with *Rhizobium* and phosphate solubilizing bacteria (*Bacillus polymyxa*) resulted in the higher rate of nitrogen fixation and phosphorus solubilization from insoluble phosphorus forms. From Table 1 it was revealed that there was significant difference in plant height due to application of *Rhizobium* and phosphate solubilizing bacteria (*Bacillus polymyxa*) seed inoculation. Highest and lowest plant height was 21.24 cm and 18.12 cm in the plots treated with inoculation (A<sub>1</sub>) and without any culture (A<sub>0</sub>), respectively at 30 DAS. The same trend was followed up to 60 DAS. The highest plant height of 23.96 cm and 52.55 cm at 30 and 60 DAS, respectively recorded in plot received the recommend dose of NPK followed by plot treated with 75 per cent recommended dose of N, P<sub>2</sub>O<sub>5</sub> and 100 per cent K<sub>2</sub>O.

Dry matter accumulations and crop growth rate (CGR) were observed. Inoculation with *Rhizobium* and phosphate solubilizing bacteria (*Bacillus polymyxa*) increased nodulation. As a result, dry matter accumulation was significantly higher (46.04 and 383.19 g m<sup>-2</sup> at 30 and 60 DAS, respectively) and also the CGR (12.69 g m<sup>-2</sup> days<sup>-1</sup>) due to seed inoculation as compared to no

inoculation (Table 1). Among the different fertilizer treatments the highest dry matter accumulation of 51.05 and 386.44 gm<sup>-2</sup> at 30 and 60 DAS, respectively was observed (Table 1) with the treatment B<sub>1</sub>, *i.e.* application of recommend dose of NPK, followed by that in the treatment B<sub>2</sub> *i.e.* application of 75 per cent recommended dose of N and P<sub>2</sub>O<sub>5</sub> + 100% K<sub>2</sub>O. Highest CGR (13.04 gm<sup>-2</sup> day<sup>-1</sup>) was observed (Table 1) with the treatment B<sub>1</sub>, *i.e.* application of recommend dose of NPK, followed by that in the treatment B<sub>2</sub> *i.e.* application of 75 per cent

recommended dose of N, P<sub>2</sub>O<sub>5</sub> and 100 per cent K<sub>2</sub>O. The lowest (37.45gm<sup>-2</sup>) dry matter accumulation was recorded in the control plot where no fertilizer was applied. The interaction effects (Table 2) between two factors were different significantly at all the growth stages. The maximum dry matter accumulation was 51.18 gm<sup>-2</sup> at 30 DAS with treatment A<sub>1</sub> x B<sub>1</sub> *i.e.* seed inoculation and application of recommended dose of NPK which was at par with A<sub>1</sub> x B<sub>2</sub> *i.e.* seed inoculation and application of 75 per cent recommended dose of N,

**Table 1: Effect of *Rhizobium* and phosphate solubilizing bacteria (main plot) with inorganic fertilizers (sub-plot) on growth parameters of greengram (mean data of two years)**

Treatments	Plant height (cm)		Dry matter accumulation (g m <sup>-2</sup> )		CGRg m <sup>-2</sup> day <sup>-1</sup>	No. of nodules plant <sup>-1</sup>			Dry weight of nodules (g plant <sup>-1</sup> )		
	30 DAS	60 DAS	30 DAS	60 DAS		30 DAS	45 DAS	60 DAS	30 DAS	45 DAS	60 DAS
<b>Biofertilizers</b>											
A <sub>0</sub>	18.12	46.49	40.67	366.41	12.46	27.62	44.43	27.62	0.023	0.047	0.0342
A <sub>1</sub>	21.24	48.34	46.04	383.19	12.69	31.50	47.66	31.46	0.028	0.059	0.0413
C.D. (P=0.05)	0.3611	1.1325	2.7021	1.7745	0.4272	0.3438	1.1677	0.7037	0.0003	0.0003	0.0011
<b>Inorganic fertilizers</b>											
B <sub>0</sub>	15.89	41.32	37.45	358.76	11.99	24.85	41.85	24.85	0.0205	0.040	0.029
B <sub>1</sub>	23.96	52.55	51.05	386.44	13.04	35.46	51.63	35.37	0.0335	0.065	0.045
B <sub>2</sub>	21.34	48.93	45.20	385.43	12.88	31.30	47.96	31.30	0.0285	0.055	0.041
B <sub>3</sub>	17.52	46.87	39.75	368.58	12.40	26.65	42.75	26.65	0.0225	0.052	0.035
C.D. (P=0.05)	0.210	1.9656	1.7572	5.5531	0.4487	0.8031	0.8682	0.5072	0.0008	0.0008	0.0014

Treatments details: Main plots- A<sub>0</sub>: no seed inoculation and A<sub>1</sub>: seed co-inoculation with PSB and *Rhizobium* (M-10) strains sub plots- B<sub>0</sub>: untreated control, B<sub>1</sub>: application of NPK (recommended dose), B<sub>2</sub>: 75% recommended dose of N and P<sub>2</sub>O<sub>5</sub> + 100% K<sub>2</sub>O, B<sub>3</sub>: 50% recommended dose of N and P<sub>2</sub>O<sub>5</sub> + 100% K<sub>2</sub>O  
DAS= Days after sowing

**Table 2: Interaction effect of *Rhizobium* and phosphate solubilizing bacteria (main plot) with a inorganic fertilizers (sub-plot) on growth parameters of greengram (mean data of two years)**

Interaction (A x B)	Plant height (cm)		Dry matter accumulation (g m <sup>-2</sup> )		CGRg m <sup>-2</sup> day <sup>-1</sup>	No. of nodules plant <sup>-1</sup>			Dry weight of nodules (g plant <sup>-1</sup> )		
	30 DAS	60 DAS	30 DAS	60 DAS		30 DAS	45 DAS	60 DAS	30 DAS	45 DAS	60 DAS
A <sub>0</sub> B <sub>0</sub>	14.16	40.06	34.58	352.00	11.94	23.90	40.84	23.90	0.019	0.033	0.027
A <sub>0</sub> B <sub>1</sub>	23.55	52.14	50.93	381.70	12.95	35.50	51.51	35.50	0.034	0.062	0.045
A <sub>0</sub> B <sub>2</sub>	18.15	45.31	39.28	371.16	12.77	26.10	43.34	26.10	0.022	0.046	0.034
A <sub>0</sub> B <sub>3</sub>	16.63	48.47	37.89	360.81	12.19	25.00	42.06	25.00	0.020	0.047	0.031
A <sub>1</sub> B <sub>0</sub>	17.62	42.58	40.32	365.53	12.04	25.80	42.87	25.80	0.022	0.048	0.031
A <sub>1</sub> B <sub>1</sub>	24.38	52.96	51.18	391.18	13.13	35.42	51.75	35.24	0.033	0.068	0.046
A <sub>1</sub> B <sub>2</sub>	24.54	52.56	51.12	399.71	12.98	36.50	52.59	36.50	0.035	0.064	0.048
A <sub>1</sub> B <sub>3</sub>	18.42	45.28	41.62	376.35	12.60	28.30	43.45	28.30	0.025	0.058	0.040
S.E.± (A x B)	0.1246	0.1311	1.0431	3.2954	0.0355	0.4767	0.5152	0.3010	0.0004	0.0005	0.0005
C.D. (P=0.05)	0.3636	0.3826	3.04	9.61	0.1036	1.3912	1.5036	0.8784	0.0011	0.0014	0.0014
S.E.± (B x A)	0.1042	0.1023	0.9342	3.0027	0.0527	0.4228	0.4849	0.2763	0.0005	0.00048	0.00046
C.D. (P=0.05)	0.3041	0.2986	2.7264	8.7554	0.1538	1.2339	1.4151	0.8063	0.0014	0.0014	0.0013

Treatments details: Main plots- A<sub>0</sub>: no seed inoculation and A<sub>1</sub>: seed co-inoculation with PSB and *Rhizobium* (M-10) strains. Sub plots- B<sub>0</sub>: untreated control, B<sub>1</sub>: application of NPK (recommended dose), B<sub>2</sub>: 75% recommended dose of N and P<sub>2</sub>O<sub>5</sub> + 100% K<sub>2</sub>O, B<sub>3</sub>: 50% recommended dose of N and P<sub>2</sub>O<sub>5</sub> + 100% K<sub>2</sub>O  
DAS= Days after sowing

$P_2O_5$  and 100 per cent  $K_2O$ .

The magnitude of nitrogen fixation by pulse crop is determined by nodule formation. The number of nodules plant<sup>-1</sup> was observed at three different stages of growth at 30, 45 and 60 DAS. There were significant differences in nodule count due to different applied fertilizers to mung bean (Table 1). The highest number of nodules per plant (31.50, 47.66 and 31.46) was found at 30, 45 and 60 DAS, respectively in the plots treated with co inoculation of seed. The lowest number of nodules per plant was recorded under the treatment  $A_0$ , i.e. no seed inoculation. Among the different fertilizer treatments the highest number of nodules plant<sup>-1</sup> was observed 35.46, 51.63 and 35.37 at 30, 45 and 60 DAS, respectively (Table 1) in the crops grown with basal application of recommended dose of NPK followed by treatment  $B_2$  i.e. application of 75 per cent recommended dose of N,  $P_2O_5$  and 100 per cent  $K_2O$ . The lowest nodules number plant<sup>-1</sup> was recorded in untreated control. Maximum number of nodule plant<sup>-1</sup> of 35.42, 51.75 and 35.24 at 30, 45 and 60 DAS, respectively were recorded in the plants grown with the treatment combination seed co-inoculation along with basal application of NPK (recommended dose) which was at par with the treatment combination of seed co-inoculation and 75 per cent recommended dose of N,  $P_2O_5$  and 100 per cent  $K_2O$  plot. In general, number of nodules plant<sup>-1</sup> increased upto first 45 DAS and, thereafter, started decreasing, irrespective of all the treatments. This is mainly due to the reduction in nutrient demand of the crop as it steps forward towards maturity.

Dry weight of nodules indicates growth of the nodule. Seed inoculation also showed the significantly the highest nodule dry weight per plant (0.0028, 0.059 and 0.0413

gm plant<sup>-1</sup> at 30, 45, and 60 DAS, respectively), while the corresponding lowest were (0.023, 0.047 and 0.034 g plant<sup>-1</sup>) in no seed inoculation. Among the different fertilizer treatment highest nodule dry weight plant<sup>-1</sup> (0.0335, 0.065 and 0.045 g plant<sup>-1</sup> at 30, 45 and 60 DAS, respectively) were recorded in recommended dose of NPK and closely followed by treatment  $B_2$  i.e. application of 75 per cent recommended dose of N and  $P_2O_5$  + 100%  $K_2O$  (Table 1). The nodule dry weight plant<sup>-1</sup> was lowest at all the growth stages with respect to the treatment with no fertilizer application ( $B_0$ ). Significant interaction effect was found at all dates of observation (Table 2), however, the best interaction effect was found with both the treatment combination  $A_1B_1$  i.e. seed inoculation with and application of recommended dose of NPK as basal (0.035, 0.064 and 0.048 g plant<sup>-1</sup>) and treatment combination  $A_1B_2$  i.e. seed inoculation and application of 75 per cent recommended dose of N and  $P_2O_5$  + 100%  $K_2O$ . The lowest (0.019, 0.033 and 0.027g plant<sup>-1</sup>) interaction effect was found with the treatment combination  $A_0B_0$  i.e. no seed inoculation and no fertilizer application. It was observed in (Table 2) that nodule dry wt. increased upto 45 DAS and then it was declined due to reduction in nodule number due to shortage of carbohydrate as the photosynthates are mostly diverted towards seed.

### Yield attributes and yield :

Number of pods plant<sup>-1</sup> and number of seed pod<sup>-1</sup> the most important yield attributes of pulses and were recorded at the time of harvesting. From the result it was found that the highest number of pods plant<sup>-1</sup> and seed pod<sup>-1</sup> was recorded (26.07 and 11.42) in the seed

**Table 3: Effect of *Rhizobium* and phosphate solubilizing bacteria (main plot) with inorganic fertilizers (sub-plot) on yield attributes of greengram (mean data of two years)**

Treatments	Pods plant <sup>-1</sup>	Seeds pod <sup>-1</sup>	Seed yield (kg ha <sup>-1</sup> )	Straw yield (kg ha <sup>-1</sup> )	Harvest index (%)
<b>Biofertilizers</b>					
$A_0$	23.695	10.9925	718.25	2484.25	22.18
$A_1$	26.077	11.4225	799.25	2457.25	22.49
C.D. (P=0.05)	1.6215	0.3069	16.65	24.6192	0.5016
<b>Inorganic fertilizers</b>					
$B_0$	20.275	9.735	632.5	2309	21.22
$B_1$	28.795	12.23	869.5	2638	23.23
$B_2$	28.17	12.255	833	2577.5	22.85
$B_3$	22.305	10.61	700	2358.5	22.04
C.D. (P=0.05)	1.4405	0.3744	16.16	25.1793	0.5799

Treatments details: Main plots- $A_0$ : no seed inoculation and  $A_1$ :seed co-inoculation with PSB and *Rhizobium* (M-10) strains. Sub plots- $B_0$ : untreated control,  $B_1$ : application of NPK (recommended dose),  $B_2$ : 75% recommended dose of N and  $P_2O_5$  + 100%  $K_2O$ ,  $B_3$ : 50% recommended dose of N and  $P_2O_5$  + 100%  $K_2O$   
DAS= Days after sowing

inoculation (Table 3). Among the different fertilizer treatments significantly the highest (28.79) number of pods plant<sup>-1</sup> and 12.23 seed pod<sup>-1</sup> were recorded in recommended dose of NPK followed by treatment B<sub>2</sub> i.e. application of 75 per cent recommended dose of N and P<sub>2</sub>O<sub>5</sub> + 100% K<sub>2</sub>O (Table 3). The maximum values of pods plant<sup>-1</sup> (30.10) were observed both under the treatment A<sub>1</sub>B<sub>1</sub> i.e. seed inoculation with application of recommended dose of NPK as basal and treatment combination A<sub>1</sub>B<sub>2</sub> i.e. seed inoculation and application of 75 per cent recommended dose of N and P<sub>2</sub>O<sub>5</sub> + 100% K<sub>2</sub>O (30.85). The maximum values of seed pod<sup>-1</sup> were observed both under the treatment A<sub>1</sub>B<sub>1</sub> i.e. seed inoculation with application of recommended dose of NPK as basal (12.27) and treatment combination A<sub>1</sub>B<sub>2</sub> i.e. seed inoculation and application of 75 per cent recommended dose of N and P<sub>2</sub>O<sub>5</sub> + 100% K<sub>2</sub>O (12.81). Singh *et al.* (2010) found similar results with lentil. Corresponding the lowest number of pods plant<sup>-1</sup> (19.86) and seed pod<sup>-1</sup> (9.52) were observed in the control. Significant interaction effect was found due to different treatment combinations (Table 4). It was found that seed co-inoculation with *Rhizobium* and phosphate solubilizing bacteria (*Bacillus polymyxa*) increased the pods plant<sup>-1</sup> as well as seed pod<sup>-1</sup> in mung bean. The increase in grain yield with nitrogen and phosphorus application might be due to higher number of pods plant<sup>-1</sup> and grains pod<sup>-1</sup> as grain yield is known to have positive association with these characters (Lavanya and Toms, 2009).

From the result it was revealed that a significant increase in seed yield and straw yield of mung bean was

found with seed co-inoculation than un-inoculated treatment (Table 3). The highest seed yield (799 kg ha<sup>-1</sup>) and straw yield (2457 kg ha<sup>-1</sup>) addressed with seed co-inoculation. This finding corroborated with the result of Yousuf *et al.* (1989). Application of recommended dose of NPK increased seed yield and straw yield significantly (Table 3) over control. Recommended dose of NPK produced the highest seed yield (869 kg ha<sup>-1</sup>). Maximum straw yield (2638 kg ha<sup>-1</sup>) was recorded against the treatment with recommended dose of NPK (B<sub>1</sub>) and lowest (2309 kg ha<sup>-1</sup>) against treatment with no fertilizer application B<sub>0</sub>. Among the different interaction combination the maximum seed yield (939 kg ha<sup>-1</sup>) and straw yield (2648 kg ha<sup>-1</sup>) was observed with A<sub>1</sub>B<sub>1</sub> i.e. seed inoculation and along with recommended dose of NPK which is at par with 927 kg ha<sup>-1</sup> and 2662 kg ha<sup>-1</sup> seed yield and straw yield, respectively, in the treatment combination A<sub>1</sub>B<sub>2</sub> i.e. seed inoculation and application of 75 per cent recommended dose of N and P<sub>2</sub>O<sub>5</sub> + 100% K<sub>2</sub>O. The lowest seed yield (649 kg ha<sup>-1</sup>) and stalk yield (2310 kg ha<sup>-1</sup>) was obtained with the treatment combination A<sub>0</sub>B<sub>0</sub> i.e. untreated control plot. Similar findings also reported by Choudhary *et al.* (2011). Significant difference was found in harvest index in different treatment combination (Table 3). The highest harvest index (22.49%) was observed in seed co-inoculation. Recommended dose of NPK showed the highest harvest index (23.23%) of mung bean. Among all the treatments significantly the highest harvest index (23.21%) was obtained in the treatment combination A<sub>1</sub>B<sub>2</sub> i.e. seed inoculation and application of 75 per cent recommended dose of N and P<sub>2</sub>O<sub>5</sub>+100% K<sub>2</sub>O (Table 4).

**Table 4: Interaction effect of *Rhizobium* and phosphate solubilizing bacteria (main plot) with inorganic fertilizers (sub-plot) on yield attributes of greengram (mean data of two years)**

Treatments	Pods plant <sup>-1</sup>	Seeds pod <sup>-1</sup>	Seed yield (kg ha <sup>-1</sup> )	Straw yield(kg ha <sup>-1</sup> )	Harvest index(%)
A <sub>0</sub> B <sub>0</sub>	19.86	9.52	649	2310	20.65
A <sub>0</sub> B <sub>1</sub>	27.49	12.19	800	2628	23.25
A <sub>0</sub> B <sub>2</sub>	25.49	11.70	739	2563	22.43
A <sub>0</sub> B <sub>3</sub>	21.94	10.56	685	2436	22.40
A <sub>1</sub> B <sub>0</sub>	20.69	9.95	616	2308	21.80
A <sub>1</sub> B <sub>1</sub>	30.10	12.27	939	2648	23.21
A <sub>1</sub> B <sub>2</sub>	30.85	12.81	927	2662	23.28
A <sub>1</sub> B <sub>3</sub>	22.67	10.66	715	2281	21.69
S.E.± (A x B)	0.8549	0.2230	7.031	14.9432	0.3441
C.D. (P=0.05)	2.4950	0.6508	20.51	43.6116	1.004
S.E.± (B x A)	0.7310	0.7202	6.272	13.622	0.3062
C.D. (P=0.05)	2.133	2.101	18.30	37.75	0.8936

Treatments details: Main plots- A<sub>0</sub>: no seed inoculation and A<sub>1</sub>: seed co-inoculation with PSB and *Rhizobium* (M-10) strains.

Sub plots- B<sub>0</sub>: untreated control, B<sub>1</sub>: application of NPK (recommended dose), B<sub>2</sub>: 75% recommended dose of N and P<sub>2</sub>O<sub>5</sub> + 100% K<sub>2</sub>O,

B<sub>3</sub>: 50% recommended dose of N and P<sub>2</sub>O<sub>5</sub> + 100 % K<sub>2</sub>O

DAS= Days after sowing

## Conclusion :

From the above experiment it was observed that inoculation with *Rhizobium* and phosphate solubilizing bacteria (PSB) along with 75 per cent RDF *i.e.* treatment combination ( $A_1B_2$ ) was at par with treatment  $A_1B_1$  *i.e.* 100 per cent RDF with respect to all the growth parameters and yield attributing characters of mungbean. So, it can be concluded that both 25 per cent nitrogenous and phosphatic fertilizer of the recommended dose can be substituted by seed co- inoculation with phosphate solubilizing bacteria and *Rhizobia* without affecting the yield compared to 100 per cent RDF.

## Literature Cited

- Algawadi, A.R. and Gaur, A.C. (1988).** Associative effect of rhizobium and phosphate solubilizing bacteria on the yield and nutrient uptake of chickpea. *Plant Soil*, **105**: 241-246.
- Ali, M. and Kumar, S. (2005).** *Yet to see a break through.* The Hindu survey of Indian Agriculture., 54-56pp.
- Chaturvedi, S.K. and Ali, M. (2002).** *Poor man's meat needs fresh fillip.* The Hindu Survey of Indian Agriculture., 63-69pp.
- Choudhary, H.R., Sharma, O.P., Yadav, L.R. and Choudhary, G.L. (2011).** Effect of organic sources and chemical fertilizers on productivity of mungbean. *J. Food Leg.*, **24**: 324- 326.
- GOI (2009). *Economic survey of India*, Ministry of Finance (Economic Division) GOI, New Delhi., 17-22 pp.
- GOWB (2006). *Economic review.* Directorate of Agriculture, Government of West Bengal., 56-58pp.
- Gupta, A., Saxena, A. K., Gopal, M. and Tilak, K.V.B. R. (1998).** Effect of plant growth promoting rhizobacteria on competitive ability of introduced *Brayrhizobium* sp. (*Vigna*) for nodulation. *Microbiol. Res.*, **153**: 113-117.
- Lavanya, G.R. and Toms, B. (2009).** Association and interrelationship among yield contributing characters in mungbean. *J. Food Leg.*, **22**: 65-67.
- Martins, A., Kimura, O., Goi, S.R. and Baldani, J. I. (2004).** Effect of co-inoculation of plant growth promoting rhizobacteria and rhizobia on development of common bean plants (*Phaseolus vulgaris* L.). *Floresta e Ambiente*, **11**: 33-39.
- Olsen, S.R., Cole, C.V., Watanabe, F.S. and Dean, L.A. (1954).** Estimation of available phosphorus in soils by extraction with sodium bicarbonate. USDA Circular 939. 19 p. United States Department of Agriculture (USDA), Washington, DC., U.S.A.
- Payasi, D., Pandey, S., Nair, S.K. and Pandey, R.L. (2011).** Genetics of biochemical and physiological parameters responsible for powdery mildew disease resistance in mungbean. *J. Food Leg.*, **24**: 292-295.
- Perveen, S., Khan, M.S. and Zaidi, A. (2002).** Effect of rhizospheric micro-organisms on growth and yield of green gram (*Phaseolus radiatus*). *Indian J. Agric. Sci.*, **72**: 421-423.
- Provorov, N.A., Sairnazarov, U.V., Bhahromy, P.D.Z., Lozhemyakov, A.P. and Kurbanov, G.S. (1998).** Effect of *Rhizobium* inoculation on the seed (herbage) production of mungbean (*Phaseolus aureus* Roxb.) grown at Uzbekistan. *J. Arid Environ.*, **39**: 569-575.
- Rasal, P.H. (1996).** Effect of phosphomicrobials and *Rhizobium* on greengram. *Madras Agric. J.*, **83**: 399-401.
- Sarkar, R.K., Karmakar, S. and Chakraborty, A. (1993).** Response of summer greengram (*Phaseolus radiatus*) to nitrogen, phosphorus application and bacterial inoculation. *Indian J. Agron.*, **38**: 578-581.
- Schachtman, D.P., Reid, R.J. and Ayling, S.M. (1998).** Phosphorus uptake by plants : From soil to cell. *Plant Physiol.*, **116**: 447-453.
- Shukla, S.K. and Dixit, R.S. (1996).** Effect of *Rhizobium* inoculation, plant population and phosphorus on growth and yield of summer greengram (*Phaseolus radiatus*). *Indian J Agron.*, **41**: 611-615.
- Singh, A.K., Choudhary, R.K. and Sharma, R.P.R. (1993).** Effects of inoculation and fertilizer levels on yield, yield attributes and nutrient uptake of greengram (*Phaseolus radiatus*) and blackgram (*P. mungo*). *Indian J. Agron.*, **38**: 663-665.
- Singh, G., Sekhon, H.S., Ram, H., Gill, K.K. and Sharma, P. (2010).** Effect of date of sowing on nodulation, growth, thermal requirement and grain yield of *Kharif* mungbean genotypes. *J. Food Leg.*, **23**: 132-134.
- Singh, G., Sekhon, H.S., Singh, G., Brar, J.S., Bains, T.S. and Shanmugasundaram, S. (2011).** Effect of plant density on the growth and yield of mungbean [*Vigna radiata* (L.) Wilczek] genotypes under different environments in India and Taiwan. *Internat. J. Agric. Res.*, **6**: 573 - 583.
- Subbiah, B.V. and Asija, G.L. (1956).** A rapid procedure for determination of available nitrogen in soils. *Curr. Sci.*, **25**: 259-260.
- Yousuf, B.R., Harcharan, S., Cheema, S.S. and Singh, H. (1989).** Effects of combined application of pesticides and fertilizers on the growth and yield of mungbean [*Vigna radiata* (L.) Wilczek]. *Indian J. Ecol.*, **8** (2): 180-188.

11<sup>th</sup>  
Year  
★★★★★ of Excellence ★★★★★